





















Structure	Cluster	ID
	1	ZINC03776439
	1	ZINC03778256
	1	ZINC03778257
	1	ZINC03834079
	1	ZINC03832329




Structure	Cluster	ID
	1	ZINC01483030
	1	ZINC03600042
	1	ZINC03788179
	1	ZINC03834085
	1	ZINC03600044

Structure	Cluster	ID
 <p>Chemical structure showing a bicyclic nucleoside derivative. The base is a fused pyrimidine-imidazole ring system with an amino group (NH₂) at the 6-position. The sugar is a five-membered ring with a fluoromethyl group (F-CH₂-) at the 2-position and two hydroxyl groups (OH) at the 3 and 4 positions. The sugar is attached to the base via a glycosidic bond.</p>	1	ZINC03782812
 <p>Chemical structure showing a bicyclic nucleoside derivative. The base is a fused pyrimidine-imidazole ring system with an amino group (NH₂) at the 6-position. The sugar is a five-membered ring with two hydroxyl groups (OH) at the 3 and 4 positions. The sugar is attached to the base via a glycosidic bond.</p>	1	ZINC03832327
 <p>Chemical structure showing a bicyclic nucleoside derivative. The base is a fused pyrimidine-imidazole ring system with an amino group (NH₂) at the 6-position. The sugar is a five-membered ring with a hydroxymethyl group (OH-CH₂-) at the 2-position and two hydroxyl groups (OH) at the 3 and 4 positions. The sugar is attached to the base via a glycosidic bond.</p>	1	ZINC03832328
 <p>Chemical structure showing a bicyclic nucleoside derivative. The base is a fused pyrimidine-imidazole ring system with an amino group (NH₂) at the 6-position. The sugar is a five-membered ring with a hydroxymethyl group (OH-CH₂-) at the 2-position and two hydroxyl groups (OH) at the 3 and 4 positions. The sugar is attached to the base via a glycosidic bond.</p>	1	ZINC03832334
 <p>Chemical structure showing a bicyclic nucleoside derivative. The base is a fused pyrimidine-imidazole ring system with an amino group (NH₂) at the 6-position. The sugar is a five-membered ring with a hydroxymethyl group (OH-CH₂-) at the 2-position and two hydroxyl groups (OH) at the 3 and 4 positions. The sugar is attached to the base via a glycosidic bond.</p>	1	ZINC03834078

Structure	Cluster	ID
	1	ZINC03834077
	1	ZINC03834082
	1	ZINC03834081
	1	ZINC03834080
	1	ZINC04628821

Structure	Cluster	ID
 <p>Chemical structure of a bicyclic nucleoside derivative. It features a fused pyrimidine-imidazole ring system with an amino group (NH₂) at the 6-position of the pyrimidine ring. The nucleoside part consists of a ribose sugar with a fluoromethyl group (-CH₂F) at the 2' position and a hydroxyl group (-OH) at the 3' position. The sugar is attached to the imidazole ring via a glycosidic bond.</p>	1	ZINC03832338
 <p>Chemical structure of a bicyclic nucleoside derivative, identical to the first entry. It features a fused pyrimidine-imidazole ring system with an amino group (NH₂) at the 6-position of the pyrimidine ring. The nucleoside part consists of a ribose sugar with a fluoromethyl group (-CH₂F) at the 2' position and a hydroxyl group (-OH) at the 3' position. The sugar is attached to the imidazole ring via a glycosidic bond.</p>	1	ZINC03832336
 <p>Chemical structure of a bicyclic nucleoside derivative. It features a fused pyrimidine-imidazole ring system with an amino group (NH₂) at the 6-position of the pyrimidine ring. The nucleoside part consists of a ribose sugar with a 2-hydroxyethyl group (-CH₂CH₂OH) at the 2' position and a hydroxyl group (-OH) at the 3' position. The sugar is attached to the imidazole ring via a glycosidic bond.</p>	1	ZINC03826694
 <p>Chemical structure of a bicyclic nucleoside derivative. It features a fused pyrimidine-imidazole ring system with an amino group (NH₂) at the 6-position of the pyrimidine ring. The nucleoside part consists of a ribose sugar with a 1-hydroxyethyl group (-CH(OH)CH₃) at the 2' position and a hydroxyl group (-OH) at the 3' position. The sugar is attached to the imidazole ring via a glycosidic bond.</p>	1	ZINC03802230
 <p>Chemical structure of a bicyclic nucleoside derivative. It features a fused pyrimidine-imidazole ring system with an amino group (NH₂) at the 6-position of the pyrimidine ring. The nucleoside part consists of a ribose sugar with a propargyl group (-CH₂C≡CH) at the 2' position and a hydroxyl group (-OH) at the 3' position. The sugar is attached to the imidazole ring via a glycosidic bond.</p>	1	ZINC03832330

Structure	Cluster	ID
	1	ZINC03832331
	1	ZINC03832332
	1	ZINC03832333
	1	ZINC03832340
	1	ZINC03832339

Structure	Cluster	ID
 <p>The structure shows a fused thiazolopyrimidine ring system. The pyrimidine ring has an amino group (NH₂) at the 6-position. The thiazole ring is substituted with a propargyl group (-CH₂-C≡CH) at the 2-position and a sulfonamide group (-SO₂NH₂) at the 4-position.</p>	1	ZINC03832341
 <p>The structure features a pyridine ring with an amino group (NH₂) at the 3-position. It is substituted at the 1-position with a 2,3,4-trihydroxypropyl group (-CH(OH)-CH(OH)-CH₂-OH).</p>	2	ZINC03834084
 <p>The structure features a pyridine ring with a methyl group (-CH₃) at the 4-position. It is substituted at the 1-position with a 2,3,4-trihydroxypropyl group (-CH(OH)-CH(OH)-CH₂-OH).</p>	2	ZINC03834083